

Properties of Two New M31 dSph Companions from Keck Imaging

Eva K. Grebel^{1,2} and Puragra Guhathakurta^{1,3,4}

¹UCO/Lick Observatory, University of California at Santa Cruz, Santa Cruz, California 95064, USA

²Hubble Fellow

³Visiting Astronomer, W. M. Keck Observatory, jointly operated by the California Institute of Technology
and the University of California

⁴Alfred P. Sloan Research Fellow

Received _____; accepted _____

ABSTRACT

We have obtained Keck Low Resolution Imaging Spectrograph images in V and I of the newly discovered Local Group dwarf galaxies Pegasus dSph and Cassiopeia dSph and their surrounding field. The first stellar luminosity functions and color-magnitude diagrams are presented for stars with $V \lesssim 25$ and $I \lesssim 24$. The distances to the new dwarfs are estimated from the apparent I mag of the tip of the red giant branch to be: $D_{\text{MW}}(\text{Peg dSph}) = 830 \pm 80$ kpc and $D_{\text{MW}}(\text{Cas dSph}) = 760 \pm 70$ kpc, consistent with their belonging to the extended M31 satellite system. Both galaxies are dwarf spheroidals (dSphs) with red giant branch morphologies indicating predominantly old stellar populations and estimated mean metallicities, $\langle [\text{Fe}/\text{H}] \rangle$, of -1.3 ± 0.3 for Peg dSph and -1.4 ± 0.3 for Cas dSph. Their central surface brightness– $\langle [\text{Fe}/\text{H}] \rangle$ –absolute magnitude relationship follows that of other Local Group dSphs, dwarf ellipticals, and dwarf irregulars. In contrast to four out of nine Milky Way dSphs (the four that lie beyond $D_{\text{MW}} = 100$ kpc), none of the six known M31 dSphs appears to have a dominant intermediate-age population.

Subject headings: Local Group – galaxies: individual [Peg dSph (And VI), Cas dSph (And VII), And V, M31] – galaxies: stellar content – galaxies: luminosity function – galaxies: photometry – galaxies: structure

1. Introduction

Recently two groups have reported the independent discovery of new Local Group dwarf galaxies, potential companions of M31. Armandroff et al. (1998a) used a digital filtering technique to survey digitized POSS-II plates covering a 1550 deg^2 area surrounding M31 and found several candidate dwarf galaxies. They demonstrated that their first candidate, And V, is a dwarf spheroidal (dSph) galaxy sufficiently close to M31 to be its satellite. An additional candidate, named And VI following van den Bergh’s (1974) convention, was resolved into stars and found not to contain HII regions (Armandroff et al. 1998b,c). The second group, Karachentsev & Karachentseva (1998a,b) visually inspected film copies of POSS-II plates and detected four potential new dwarf companions of M31. The two brightest, granulated detections were deemed to be the most promising candidates. These were initially named PegasusDw and CassiopeiaDw after their parent constellations (Karachentsev & Karachentseva 1998a,b), and later renamed Peg dSph and Cas dSph by Tikhonov & Karachentsev 1999). Note, Peg dSph is equivalent to And VI and was discovered independently by the two groups. The galaxy Cas dSph is located in an area for which no POSS-II plate scan was available to Armandroff et al. (1998a). We will adopt the designations Peg dSph and Cas dSph throughout this *Letter*.

These three newly discovered candidate M31 dSphs, And V, Peg dSph, and CasDw, double the census of known M31 dSphs. It is important to test whether the M31 dSphs follow the same relations between absolute magnitude, surface brightness, metallicity, M/L ratio, and galactocentric distance as the Milky Way dSph satellites? Do environment and distance from M31 affect their formation histories as one might expect from van den Bergh’s (1994) ram pressure/tidal stripping scenario?

This *Letter* presents results from Keck imaging of Peg dSph and Cas dSph that address some of these questions; a preliminary version of the results were presented by Grebel (1999). Our Keck results are also compared to the results from three other recent/ongoing studies of a similar nature targeting these galaxies (Tikhonov & Karachentsev 1999; Hopp et al. 1999; Armandroff et al. 1998c, 1999).

2. Observations and Data Reduction

In preparation for a stellar spectroscopy program of M31’s dSph companions, Peg dSph and Cas dSph were imaged with the Keck II 10-m telescope in August–September 1998, using the Low Resolution Imaging Spectrograph (Oke et al. 1995) with a 2048^2 CCD and a scale of $0''.215 \text{ pixel}^{-1}$. Short exposures were

obtained in the V and I bands: 1×300 s and 3×200 s, respectively, per galaxy. Observing conditions were photometric and the seeing ranged from $0''.7$ – $1''.0$ (FWHM). The overlapping field of view of the final, combined images is $4'.6 \times 7'.1$. Resolved objects (galaxies) and regions around bright, saturated stars are excluded from further analysis (cf. Reitzel et al. 1998). Stellar photometry is carried out with the DAOPHOT/ALLFRAME programs (Stetson 1994). The photometric transformation uses airmass coefficients from Krisciunas et al. (1987) and observations of faint secondary standards (Stetson, priv. comm.) in Landolt (1992) fields.

3. Morphological Appearance

PegdSph and CasdSph have the appearance of typical low-surface brightness spheroidal galaxies without recognizable concentrations of bright stars, H II regions, or obvious globular clusters (Figure 1). PegdSph is not detected in H α , indicating the absence of currently star-forming regions (Armandroff 1998a). An ellipticity of 0.3 is estimated for both galaxies, along with position angles of $160^\circ \pm 20^\circ$ and $55^\circ \pm 20^\circ$ for PegdSph and CasdSph, respectively. Hopp et al.’s (1999) measurement of the PegdSph structural parameters is in agreement with this estimate. The ellipticities of PegdSph and CasdSph are similar to those of Galactic dSph satellites at larger distances from the Milky Way (e.g., Caldwell et al. 1992), suggesting little or no distortion through tidal effects. A detailed analysis of the galaxies’ structural parameters will be presented in a later paper (Guhathakurta & Grebel 1999).

4. Stellar Luminosity Functions and Distances

The cumulative stellar I -band luminosity function (LF) for both galaxies is shown in Figure 2. The LF of CasdSph rises at $I \approx 20.7 \pm 0.1$, and this is interpreted as the tip of the red giant branch (TRGB). The LF upturn is more gradual for PegdSph with $I_{\text{TRGB}} \approx 20.7 \pm 0.1$. The Galactic line-of-sight reddening is estimated from DIRBE/IRAS maps (Schlegel et al. 1998) to be $E_{B-V} = 0.04 \pm 0.01$ and 0.17 ± 0.03 , or $A_I = 0.09 \pm 0.01$ and 0.35 ± 0.06 , for PegdSph and CasdSph, respectively. An absolute magnitude of $M_I = -4$ for the TRGB (Lee et al. 1993) implies true distance moduli of $(m - M)_0 = 24.6 \pm 0.2$ and 24.4 ± 0.2 for PegdSph and CasdSph, respectively, where the errors include uncertainties in the exact location of the TRGB (Poisson error, possible presence of intermediate-age stars brighter than the TRGB), reddening, and photometric calibration.

Tikhonov & Karachentsev (1999) find apparent distance moduli for Peg dSph and Cas dSph that translate into true distance moduli of $(m - M)_0 = 24.52$ and 24.36 , respectively, when the above foreground extinction correction is applied. Corrected in the same manner, Hopp et al.’s (1999) finding corresponds to $(m - M)_0 = 24.43 \pm 0.2$ for Peg dSph. These results are in excellent agreement with the results of this study, when the uncertainties in the measurements are taken into account.

5. Color-Magnitude Diagrams and Metallicity Estimates

The color-magnitude diagrams (CMDs) of Peg dSph and Cas dSph show prominent red giant branches (RGBs) and no evidence for young blue stars (Figure 3). Note, the surrounding $R > 3'$ “field” CMDs in Figure 3, each drawn from a region whose area is about 45% that of the $R < 3'$ galaxy area. The field CMDs are slightly contaminated by stars in the outskirts of the dSph galaxies ($< 10\%$ of the number of stars in the corresponding $R < 3'$ galaxy CMD), but this is unimportant for the purposes of this study since no significant radial gradient appears to be present in the galaxies’ properties. The Cas dSph RGB is more densely populated than that of Peg dSph, a reflection of Cas dSph’s higher stellar surface density. A small metal-rich ($[\text{Fe}/\text{H}] < -1$) population appears to be indicated by stars with $V - I \geq 2$ near the TRGB in Cas dSph (Figure 3). The density and color distribution of stars above the Cas dSph TRGB are similar to that in the corresponding “field” CMD, but the presence of a small intermediate-age asymptotic giant branch (AGB) population cannot be ruled out. The Peg dSph CMD displays an excess of stars above the TRGB relative to the field CMD; this is also seen as a tail in the LF brighter than the TRGB. These super-TRGB stars may indicate the presence of a significant intermediate-age AGB population in Peg dSph; they are not likely to be blend artifacts (cf. Grillmair et al. 1996; Martínez-Delgado & Aparicio 1997).

The mean metallicity is computed from the median reddening-corrected color, $(V - I)_0$, at $M_I = -3.5$ (Armandroff et al. 1993; Caldwell et al. 1998), computed over the range $1 < (V - I)_0 < 2$. Outside this color range the surface density of objects is identical for the inner $R < 3'$ and surrounding $R > 3'$ regions, indicating that the majority of these objects are foreground or background sources. For Peg dSph, $(V - I)_{0,-3.5} = 1.46 \pm 0.02$ for $21.1 \leq I \leq 21.3$, which yields $\langle [\text{Fe}/\text{H}] \rangle = -1.35 \pm 0.07$. For Cas dSph, $(V - I)_{0,-3.5} = 1.43 \pm 0.01$ for $21.1 \leq I \leq 21.3$, which yields $\langle [\text{Fe}/\text{H}] \rangle = -1.42 \pm 0.04$. These errors are the formal errors of the mean; the overall uncertainty in the mean metallicities is conservatively estimated to be ± 0.3 dex due to systematic errors in photometry/calibration. These metallicity estimates are in good agreement with those derived from the standard globular cluster RGB fiducials (Da Costa & Armandroff

1990) shown in Figure 3.

The observed width of the RGB in the Peg dSph and Cas dSph CMDs, $\delta(V - I) \approx 0.4$, corresponds to a metallicity spread of roughly 1 dex. The formal uncertainty in the color measurement (see Fig. 3) is significantly smaller than the observed spread in bright RGB color. The effect of systematic errors (e.g., crowding, flat fielding), however, may be as large as $\sigma(V - I) \sim 0.1$, judging from a comparative analysis of a subset of the data obtained in poorer seeing and at a different CCD orientation with respect to the primary data set. The fact that the RGB widths are similar for Peg dSph and Cas dSph despite the substantial difference in their stellar surface density argues against the widths being caused by crowding-related photometric error. Finally, the apparent flaring of the Cas dSph RGB near its bright tip is strongly suggestive of a spread in $[\text{Fe}/\text{H}]$.

Measurements of the mean metallicity in three radial bins ($R < 1'$, $1' < R < 2'$, and $2' < R < 3'$), based on a wider I -magnitude range ($\Delta I = 0.5$ mag), places a robust upper limit on any radial metallicity gradient in the galaxies. The variation in median $(V - I)_0$ color over the inner $3'$ region is found to be < 0.03 mag corresponding to a variation in mean metallicity of < 0.05 dex, comparable to the formal error in $\langle [\text{Fe}/\text{H}] \rangle$ quoted above. Thus Peg dSph and Cas dSph show no evidence for a radial metallicity gradient, in contrast with what is seen in some dSphs with substantial intermediate-age populations (see Grebel 1997 for a summary).

6. Central Surface Brightness

The central surface brightness in the V band, measured in a $40''$ -diameter aperture after subtraction of bright foreground stars and corrected for line-of-sight extinction (same procedure as in Armandroff et al. 1998a), is $\mu_{V,0} \approx 24.5$ and 23.6 mag arcsec $^{-2}$ for Peg dSph and Cas dSph, respectively; these brightnesses are higher than the observed $\mu_{V,0}$ values for the other M31 dSphs (Caldwell et al. 1992; Armandroff et al. 1998a): 24.9 (And I), 24.8 (And II), 25.3 (And III), and 25.2 mag arcsec $^{-2}$ (And V). Peg dSph and Cas dSph, along with the newly discovered And V dSph (Armandroff et al. 1998a), follow the general $\langle [\text{Fe}/\text{H}] \rangle$ -vs- $\mu_{V,0}$ relationship defined by Local Group dwarf ellipticals (dEs), dwarf irregulars, and dSphs (Figure 4, *upper panel*). Absolute B -band magnitudes, M_B , are computed from the total apparent magnitudes, B_T , estimated by Karachentsev & Karachentseva (1998a). Peg dSph lies on the mean M_B -vs- $\mu_{V,0}$ relation defined by other Local Group dwarf galaxies (Figure 4, *lower panel*), while Cas dSph is somewhat less luminous than other dwarfs of comparable $\mu_{V,0}$.

7. Discussion

A summary of the properties of the two new dSphs PegdSph and CasdSph derived from Keck imaging is given in Table 1. Their mean metallicities resemble that of the M31 globular cluster system ($[\text{Fe}/\text{H}] = -1.2$ —Huchra et al. 1991) and of the intermediate-age/old field populations in M31’s dE companions, but are higher than the $\langle[\text{Fe}/\text{H}]\rangle$ of the other four M31 dSphs (cf. Grebel 1997; Armandroff et al. 1998a). PegdSph and CasdSph also have the highest central surface brightnesses of the known M31 dSphs. Their detection may have been hampered by the presence of nearby, bright foreground stars.

Heliocentric distances of 830 ± 80 kpc for PegdSph and 760 ± 70 kpc for CasdSph translate into distances of 280 ± 85 kpc and 215 ± 75 kpc from M31, respectively, adopting the Cepheid-based distance of 770 ± 30 kpc to M31 (cf. Freedman & Madore 1990; Holland 1998). Thus PegdSph and CasdSph are the most distant of the currently known M31 dSph satellites. The Leo I and Leo II dSphs have comparable distances from the Milky Way, lending credence to Karachentsev & Karachentseva’s (1998a) suggestion that PegdSph and CasdSph belong to an extended subsystem of M31 companions together with And V and LGS 3.

The fraction of stars belonging to an intermediate-age population shows a bimodal distribution among the Milky Way dSphs, with small intermediate-age fractions at small Galactocentric distances and dominant fractions at large distances (van den Bergh 1994). By contrast, all the M31 dSph companions appear to have small intermediate-age populations (with the possible exception of And II—Da Costa 1998), even though they are located at comparable distances from their more massive parent galaxy. The possibility of highly eccentric orbits in the M31/M33 system cannot be ruled out. Our planned spectroscopic observations will help constrain the orbits of the M31 dSphs through radial velocity measurements and should lead to more accurate metallicity and abundance spread determinations.

We thank Igor Karachentsev, Valentina Karachentseva, Taft Armandroff, and George Jacoby for helpful discussions, and an anonymous referee for useful comments. Mike Bolte kindly provided IRAF and DAOPHOT scripts for reduction of the Keck data. We are indebted to Peter Stetson for making available his secondary standards, without which a reliable calibration would not have been possible. EKG gratefully acknowledges support by Dennis Zaritsky through NASA LTSA grant NAG-5-3501 and by NASA through grant HF-01108.01-98A from the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS5-26555.

REFERENCES

- Armandroff, T. E., Da Costa, G. S., Caldwell, N., & Seitzer, P. 1993, *AJ*, 106, 986
- Armandroff, T. E., Davies, J. E., & Jacoby, G. H. 1998a, *AJ*, 116, 2287
- Armandroff, T. E., Davies, J. E., & Jacoby, G. H. 1998b, in *Dwarf Tales*, eds. E. Brinks & E. K. Grebel, Vol. 3, p. 2
- Armandroff, T. E., Davies, J. E., & Jacoby, G. H. 1998c, in *IAU Coll. 171, The Low Surface Brightness Universe*, eds. J. I. Davies, C. Impey, & S. Philipps (San Francisco: ASP), in press
- Armandroff, T. E., Davies, J. E., & Jacoby, G. H. 1999, in preparation
- Caldwell, N., Armandroff, T. E., Seitzer, P., & Da Costa, G. S. 1992, *AJ*, 103, 840
- Caldwell, N., Armandroff, T. E., Da Costa, G. S., & Seitzer, P. 1998, *AJ*, 115, 535
- Da Costa, G. S., & Armandroff, T. E. 1990, *AJ*, 100, 162
- Da Costa, G. S. 1998, in *Stellar Astrophysics for the Local Group: A First Step to the Universe*, eds. A. Aparicio & A. Herrero (Cambridge: Cambridge University Press), in press
- Freedman, W. L., & Madore, B. F. 1990, *ApJ*, 365, 186
- Grebel, E. K. 1997, *Reviews in Mod. Astron.*, 10, 29
- Grebel, E. K. 1999, in *IAU Symp. 192, The Stellar Content of the Local Group*, eds. P. Whitelock & R. Cannon (San Francisco: ASP), in press
- Grillmair, C. J., Lauer, T. R., Worthey, G., et al. 1996, *AJ*, 112, 1975
- Guhathakurta, P., & Grebel, E. K. 1999, *ApJ*, in preparation
- Holland, S. 1998, *AJ*, 115, 1916
- Hopp, U., Schulte-Ladbeck, R. E., Greggio, L., & Mehlert, D. 1999, *A&AL*, submitted
- Huchra, J. P., Kent, S. M., & Brodie, J. P. 1991, *ApJ*, 370, 495

Karachentsev, I. D., & Karachentseva, V. E. 1998a, in Dwarf Tales, eds. E. Brinks & E. K. Grebel, Vol. 3, p.

1

Karachentsev, I. D., & Karachentseva, V. E. 1998b, A&A, submitted

Krisciunas, K., Sinton, W., Tholen, K., et al. 1987, PASP, 99, 887

Landolt, A. U. 1992, AJ, 104, 340

Lee, M. G., Freedman, W. L., & Madore, B. F. 1993, ApJ, 417, 553

Mateo, M. 1998, ARA&A, 36, 435

Martínez-Delgado, D., & Aparicio, A. 1997, ApJ, 480, 107

Oke, J. B., Cohen, J. G., Carr, M., et al. 1995, PASP, 107, 375

Reitzel, D. B., Guhathakurta, P., & Gould, A. 1998, AJ, 116, 707

Schlegel, D. J., Finkbeiner, D. P., & Davis, M. 1998, ApJ, 500, 525

Stetson, P. B. 1994, PASP, 106, 250

Tikhonov, N. A., & Karachentsev, I. D. 1999, AstL, submitted

van den Bergh, S. 1974, ApJ, 191, 271

van den Bergh, S. 1994, AJ, 108, 2145

Fig. 1.— Negative grayscale representation of V -band images of Pegasus dSph and Cassiopeia dSph (1×300 s for each galaxy) obtained with the Low Resolution Imaging Spectrograph on the 10-m Keck II telescope in August 1998. The seeing FWHM for these observations was $0''.7\text{--}1''.0$.

Fig. 2.— Cumulative I -band luminosity functions (LFs) of the central $R < 3'$ regions of Peg dSph and Cas dSph and of the surrounding $R > 3'$ field regions (scaled to match the area of the inner $R < 3'$ region). The arrows mark the adopted position of the tip of the red giant branch (TRGB) in the field-star subtracted LF (dotted line). The bright end of the field-star subtracted LF appears to extend to slightly brighter magnitudes than the adopted TRGB location, especially for Peg dSph; this may indicate the presence of luminous AGB stars.

Fig. 3.— Color-magnitude diagrams (CMDs) of the inner $R < 3'$ region of Peg dSph and Cas dSph, with their corresponding surrounding $R > 3'$ field regions (each field region has roughly 45% the area of the inner $R < 3'$ region). The error bands in the right panels represent the formal (DAOPHOT) $\pm 1\sigma$ error in $V - I$ color; the overall errors are likely to be somewhat larger. Note, the “field” CMD is slightly contaminated by red giant stars in the outer regions (> 60 pc) of the dSphs. The solid grey lines in the left panels are globular cluster fiducials from Da Costa & Armandroff (1990), appropriately reddened and shifted to the distance moduli of the galaxies, for a range of $[\text{Fe}/\text{H}]$ values: (left to right) M15 (-2.2), NGC 6752 (-1.5), NGC 1851 (-1.2), and 47 Tuc (-0.7).

Fig. 4.— (a) Mean metallicity, $\langle [\text{Fe}/\text{H}] \rangle$, plotted versus central surface brightness in the V band, $\mu_{V,0}$, for nearby dwarf galaxies. The data are from Grebel (1997), Armandroff et al. (1998a), and the present study. Peg dSph and Cas dSph (two circled dots) follow the mean $\langle [\text{Fe}/\text{H}] \rangle$ -vs- $\mu_{V,0}$ relation within the errors. (b) Absolute B -band magnitude, M_B , versus central surface brightness for the same galaxies that are plotted in (a) with the exception of the And V dSph for which an M_B measurement is unavailable. The M_B values are uncertain at the level of ± 0.5 mag (Karachentsev, priv. comm.). The data are from Mateo (1998) and the present study.

Table 1. Properties of Peg dSph and Cas dSph


Parameter	Peg dSph	Cas dSph
α (J2000)	23 ^h 51 ^m 39 ^s	23 ^h 26 ^m 31 ^s
δ (J2000)	24°35′42″	50°41′31″
l, b	106.0°, −36.3°	109.5°, −9.9°
Position angle	160° ± 20°	55° ± 20°
Ellipticity	0.3 ± 0.1	0.3 ± 0.1
E_{B-V}	0.04 ± 0.01	0.17 ± 0.03
$(m - M)_0$	24.6 ± 0.2	24.4 ± 0.2
Distance _{MW}	830 ± 80 kpc	760 ± 70 kpc
Distance _{M31}	280 ± 85 kpc	215 ± 75 kpc
Distance _{M33}	340 ± 90 kpc	430 ± 80 kpc
B_T	14.5 ± 0.5 ^a	16.0 ± 0.5 ^a
V_T	14.1 ± 0.2 ^b	
$\mu_{V,0}$	24.5 ± 0.2 mag arcsec ^{−2}	23.6 ± 0.2 mag arcsec ^{−2}
$\langle[\text{Fe}/\text{H}]\rangle$	−1.3 ± 0.3	−1.4 ± 0.3

^aFrom Karachentsev & Karachentseva (1998).

^bFrom Hopp et al. (1999), corrected using Schlegel et al.’s (1998) extinction estimate.

This figure "fig1.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/9812086v1>



This figure "fig3.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/9812086v1>

